

Parent Name	INSTRUCTIONS:	This form for any child in grades Complete and sign form. Submit	K – 12 who is unable to receive a va ted to school as proof of exemption t	accine required From required in	for school entry due to a medical contraindication. nmunization.
Parent/Guardian Name	Patient Name			Date of E	Birth (month/dav/vear)
Street Address					
City					
General Contraindications to All Vaccines (Vaccine should not be given.)					
Vaccine Specific Contraindications (Vaccine should not be given.) DTaP or Tdap	Severe allergic re Hepatitis B (Diphtheria, t	eaction (e.g., anaphylaxis) after a Hep B) etanus, pertussis (DTaP, Tdap)	previous vaccine dose or to a v. Inactivated poliovirus (IPV) Measles, mumps, rubella (N	en.) accine compo	onent Meningococcal, conjugate (MCV4) or Meningococcal, polysaccharide
Vaccine Specific Contraindications (Vaccine should not be given.) DTaP or Tdap	Which vaccine or	vaccine component caused read	ction?		
Vaccine Specific Contraindications (Vaccine should not be given.)					
DTaP or Tdap					
Known severe immunosdeficiency (e.g., hematologic and solid tumors; receiving chemotherapy; congenital immunosdeficiency; long term immunosuppressive therapy; or patients with HIV infection who are severely immunocompromised) Varcinella Pregnancy Estimated Date of Confinement (EDC):		☐ Encephalopathy (e.g., coma, d within seven (7) days of admin	ecreased level of consciousness, pro- istration of previous dose of DTP or	DTaP	
Varicella Pregnancy Estimated Date of Confinement (EDC):	MMR	☐ Pregnancy Estimated Date of	of Confinement (EDC):		(month, day year)
Pregnancy Estimated Date of Confinement (EDC):		term immunosuppressive thera	cy (e.g., hematologic and solid tumo apy; or patients with HtV infection wh	rs; receiving ch	emotherapy; congenital immunodeficiency; long immunocomoromised)
Vaccine Specific Precautions (Vaccine may be given or held depending on clinical situation.) DTaP or Tdap Guillan-Barre syndrome (GBS) within six (6) weeks after a previous dose of tetanus-containing vaccine History of Arthus-type hypersensitivity reaction following a previous dose of tetanus and/or diphtheria toxoid-containing vaccine: defer vaccination until at least ten (10) years have elapsed since the previous dose of tetanus and/or diphtheria toxoid-containing vaccine: defer vaccination until at least ten (10) years have elapsed since the previous dose of Progressive or unstable neurologic disorder, uncontrolled seizures or progressive encephalopathy: defer vaccination with DTaP or Tdap until a treatment regiment has been established and the condition has stabilized DTaP Temperature of ≥105F (≥40.5C) within forty-eight (48) hours after vaccination with a previous dose of DTP/DTaP Collapse and shock-like state (i.e.: hypotonic hyporesponsive episode) within forty-eight (48) hours after previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) days after receiving a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/	Varicella	☐ Pregnancy Estimated Date of	of Confinement (EDC):		(month, day year)
DTaP or Tdap Guillan-Barre syndrome (GBS) within six (6) weeks after a previous dose of tetanus-containing vaccine History of Arthus-type hypersensitivity reaction following a previous dose of tetanus and/or diphtheria toxoid-containing vaccine: defer vaccination until at least ten (10) years have elapsed since the previous dose Progressive or unstable neurologic disorder, uncontrolled seizures or progressive encephalopathy: defer vaccination with DTaP or Tdap until a freatment regiment has been established and the condition has stabilized Temperature of ≥105F (≥40.5C) within forty-elight (48) hours after vaccinom with a previous dose of DTP/DTaP Collapse and shock-like state (i.e.: hypotonic hyporesponsive episode) within forty-elight (48) hours after previous dose of DTP/DTaP Seizure or convulsion within three (3) days after receiving a previous dose of DTP/DTaP Seizure or convulsion within three (3) days after receiving a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-elight (48) hours after a previous dose of DTP/DTaP Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) History of thrombocytopenia or thrombocytopenic purpura Varicella Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) Receipt of specific antivirals (i.e., acyclovir, famciclovir, or valacyclovir) twenty-four (24) hours before vaccination; if possible, delay resumption of these antiviral drugs for fourteen (14) days after vaccination Other Medical Contraindication (Must list vaccine(s) and contraindications individually – continue on back if necessary.) Vaccine Specific Contraindication Office Address Please indicate the duration of the medical exemption, and if and when vaccine can be safely administered. (Exemption can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption is temporary (<1 year,) and resolu		Substantial suppression of cells	ular immunity		
DTaP or Tdap Guillan-Barre syndrome (GBS) within six (6) weeks after a previous dose of letanus-containing vaccine History of Arthus-type hypersensitivity reaction following a previous dose of tetanus and/or diphtheria toxoid-containing vaccine: defer vaccination until at least ten (10) years have elapsed since the previous dose Progressive or unstable neurologic disorder, uncontrolled seizures or progressive encephalopathy; defer vaccination with DTaP or Tdap until a treatment regiment has been established and the condition has stabilized Temperature of ≥105F (≥40.5C) within forty-elight (48) hours after vaccination with a previous dose of DTP/DTaP Collapse and shock-like state (i.e.; hypotonic hyporesponsive episode) within forty-elight (48) hours after previous dose of DTP/DTaP Seizure or convulsion within three (3) days after receiving a previous dose of DTP/DTaP Seizure or convulsion within three (3) days after receiving a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-elight (48) hours after a previous dose of DTP/DTaP Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) History of thrombocytopenia or thrombocytopenic purpura Varicella Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) Receipt of specific antivirals (i.e., acyclovir, famciclovir, or valacyclovir) twenty-four (24) hours before vaccination; if possible, delay resumption of these antiviral drugs for fourteen (14) days after vaccination Other Medical Contraindication (Must list vaccine(s) and contraindications individually – continue on back if necessary.) Vaccine Specific Contraindication Office Address Please indicate the duration of the medical exemption, and if and when vaccine can be safely administered. (Exemption can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption is temporary (<1 year), and res	Vaccine Specif	ic Precautions (Vaccine mag	y be given or held dependina	on clinical s	ituation \
History of Arthus-type hypersensitivity reaction following a previous dose of tetanus and/or diphtheria toxoid-containing vaccine:		Guillan-Barre syndrome (GBS)	b) within six (6) weeks after a previou	s dose of tetan	us-containing vaccine
DTaP		 History of Arthus-type hypersed defer vaccination until at least 	ensitivity reaction following a previou ten (10) years have elapsed since t	s dose of tetani he previous dos	us and/or diphtheria toxoid-containing vaccine:
Collapse and shock-like state (i.e.: hypotonic hyporesponsive episode) within forty-eight (48) hours after previous dose of DTP/DTaP Seizure or convulsion within three (3) days after receiving a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Presistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Presistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Presistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Presistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Presistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Presistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Presistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Presistent, inconsolate a previous dose of DTP/DTaP Presistent		 Progressive or unstable neuro Tdap until a treatment regime 	plogic disorder, uncontrolled seizures int has been established and the con	or progressive dition has stabi	encephalopathy: defer vaccination with DTaP or lized
Seizure or convulsion within three (3) days after receiving a previous dose of DTP/DTaP Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) History of thrombocytopenia or thrombocytopenic purpura Varicetla Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) Receipt of specific antivirals (i.e., acyclovir, famciclovir, or valacyclovir) twenty-four (24) hours before vaccination; if possible, delay resumption of these antiviral drugs for fourteen (14) days after vaccination Other Medical Contraindication (Must list vaccine(s) and contraindications individually – continue on back if necessary.) Vaccine Specific Contraindication Please indicate the duration of the medical exemption, and if and when vaccine can be safely administered. (Exemption can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.) Medical exemption is permanent, and will apply for one (1) year from today's date. Medical exemption is temporary (<1 year), and resolution is anticipated by	DTaP	☐ Temperature of ≥105F (≥40.5	C) within forty-eight (48) hours after	vaccination with	a previous dose of DTP/DTaP
Persistent, inconsolable crying lasting three (3) or more hours within forty-eight (48) hours after a previous dose of DTP/DTaP MMR		☐ Collapse and shock-like state ☐ Seizure or consulsion within the	(i.e.: hypotonic hyporesponsive epis	ode) within forty	y-eight (48) hours after previous dose of DTP/DTaP
Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) History of thrombocytopenia or thrombocytopenia purpura Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) Receipt of specific antivirals (i.e., acyclovir, famciclovir, or valacyclovir) twenty-four (24) hours before vaccination; if possible, delay resumption of these antiviral drugs for fourteen (14) days after vaccination Other Medical Contraindication (Must list vaccine(s) and contraindications individually – continue on back if necessary.) Vaccine Specific Contraindication Please indicate the duration of the medical exemption, and if and when vaccine can be safely administered. (Exemption can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.) Medical exemption is permanent, and will apply for one (1) year from today's date. Medical exemption is temporary (<1 year), and resolution is anticipated by/_ Medical exemption is pregnancy, and Estimated Date of Confinement (EDC) is/_/ Physician Name Office Address Telephone	•	☐ Persistent, inconsolable crying	lasting three (3) or more hours with	us aose of DIF in forty-eight (4	/DISP
History of thrombocytopenia or thrombocytopenic purpura Recent (within eleven (11) months) receipt of antibody-containing blood product (interval depends on product) Receipt of specific antivirals (i.e., acyclovir, famciclovir, or valacyclovir) twenty-four (24) hours before vaccination; if possible, delay resumption of these antiviral drugs for fourteen (14) days after vaccination Other Medical Contraindication (Must list vaccine(s) and contraindications individually – continue on back if necessary.) Vaccine Specific Contraindication Please indicate the duration of the medical exemption, and if and when vaccine can be safely administered. (Exemption can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.) Medical exemption is permanent, and will apply for one (1) year from today's date. Medical exemption is temporary (<1 year), and resolution is anticipated by	MMR	Recent (within eleven (11) mo	inths) receipt of antibody-containing	blood product (i	interval depends on product)
Receipt of specific antivirals (i.e., acyclovir, famciclovir, or valacyclovir) twenty-four (24) hours before vaccination; if possible, delay resumption of these antiviral drugs for fourteen (14) days after vaccination Other Medical Contraindication (Must list vaccine(s) and contraindications individually – continue on back if necessary.) Vaccine Specific Contraindication Please indicate the duration of the medical exemption, and if and when vaccine can be safely administered. (Exemption can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.) Medical exemption is permanent, and will apply for one (1) year from today's date. Medical exemption is temporary (<1 year), and resolution is anticipated by // / Medical exemption is pregnancy, and Estimated Date of Confinement (EDC) is // / Physician Name Physician License Number Telephone		History of thrombocytopenia of	r thrombocytopenic purpura		· · · · · · · · · · · · · · · · · ·
Please indicate the duration of the medical exemption, and if and when vaccine can be safely administered. (Exemption can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.) Medical exemption is permanent, and will apply for one (1) year from today's date. Medical exemption is temporary (<1 year), and resolution is anticipated by// Medical exemption is pregnancy, and Estimated Date of Confinement (EDC) is// Physician Name	Varicella	 Receipt of specific antivirals (i 	e., acyclovir, famciclovir, or valacycl	ovirl twenty-for	interval depends on product) Ir (24) hours before vaccination; if possible, delay
Please indicate the duration of the medical exemption, and if and when vaccine can be safely administered. (Exemption can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.) Medical exemption is permanent, and will apply for one (1) year from today's date. Medical exemption is temporary (<1 year), and resolution is anticipated by// Medical exemption is pregnancy, and Estimated Date of Confinement (EDC) is/_/ Physician Name	Other Medical (Contraindication (Must list va	accine(s) and contraindication	ns individuali	ly - continue on back if necessary \
Cateription can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.)	Vaccine	Specific Contraindication			2 and on Dear in Hoodestary;
Cateription can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.)					
Cateription can last for a maximum of one (1) year, and a new form must be completed annually if medical exemption still applies.)					
Office Address Telephone	☐ Medical e	extern a maximum of one (1) year exemption is permanent, and will exemption is temporary (<1 year) exemption is pregnancy, and Esti	, and a new form must be completed apply for one (1) year from todated, and resolution is anticipated by mated Date of Confinement (ED)	leted annually y's date. C) is/	rif medical exemption still applies.)

Dear	Parent	/Guardian:
------	--------	------------

Your child has a medical/religious exemption to vaccination and is not fully immunized. Although your child remains at risk for getting a vaccine preventable disease, IC 20-34-4 permits your child to attend school.

In the event of an outbreak of a vaccine preventable disease for which your child is not fully vaccinated, your child may be excluded from school to protect his/her health and the health of all our students and staff. It is important to understand that with some diseases such as measles, one infected child is an outbreak. The length of time your child will be kept out of school depends on the disease. Your child's exclusion may be as long as 3-4 weeks.

If your child is excluded from school, your child will also be excluded from school sponsored activities, such as sporting events, dances, and graduation that occur within the exclusion period. The school will notify you when your child can return to school.

Incompletely vaccinated children can be excluded from school due to cases of measles, chickenpox, pertussis, mumps, or any other vaccine preventable disease (at the discretion of the local health officer).

Acknowledgement of Consequences of Incomplete Vaccination

I understand that my child may be excluded from school in the event of an outbreak of a vaccine preventable disease.

I understand that school exclusion includes after-school activities, such as sporting events, dances, and graduation.

I understand that my child may be required to stay home for multiple weeks during an outbreak of a vaccine preventable disease for which he/she is not vaccinated.

Parent's name	40.000	
Signature	Date	
Child's name		

IMPORTANT NOTE: This form must be signed by a parent on this side and a physician on the other side to be valid.